Obesity & BMI Coding Tips



BMI can be misleading without an associated diagnosis because the calculation does not account for muscle mass, bone density, body composition, or ethnic or gender differences¹.

Adult BMI Weight Status

| Adult BMI (Ag | ge 20+) |
|---|-------------------|
| BMI < 19.9 | Underweight |
| BMI 20 - 24.9 | Healthy Weight |
| BMI 25 - 29.9 | Overweight |
| BMI 30 - 34.9 | Obese |
| BMI 35 - 39.9 (With 1+ Co-Morbidities) BMI ≥ 40 | 40 Morbidly Obese |

Calculating Pediatric BMI (Age 2 - 19)

Birth to 38 Months

LENGTH + WEIGHT FOR AGE HEAD CIRCUMFERENCE-FOR-AGE + WEIGHT FOR LENGTH

2 to 20 years

STATURE-FOR-AGE + WEIGHT-FOR-LENGTH BMI-FOR-AGE **OR**

2 to 5 years WEIGHT-FOR-STATURE

| Percentile | ICD-10 |
|----------------------|--------|
| < 5.0 | Z68.51 |
| 5.0 - 84.99 | Z38.52 |
| 85.0 - 94.99 - 84.99 | Z68.53 |
| ≥ 95.0 | Z68.54 |

| ICD-10 Official Guidelines ² | Document | Remember |
|---|--|--|
| BMI may be documented by any clinician. The treating provider must be the one to document obesity, morbid obesity, or any other diagnosis-related code from a BMI measurement. | Documentation must include: • Date of exam • Weight • BMI | State the clinical significance of obesity on the patient's health. Possible complications and co-morbidities include ⁴ : • Diabetes • Disorder of lipid proteins |
| BMI codes should only be reported as a secondary diagnosis code. | Documentation should include specific details of the cause and | Heart diseaseRespiratory problemsOsteoarthritis |
| BMI has no risk value without an associated diagnosis, such as obesity. ³ | extent of obesity like: • Due to excess calories | Individuals who are overweight, obese or morbidly obese are at an increased risk for certain medical conditions when compared |
| Coders and billers cannot infer obesity from a BMI value. | Endocrine relatedMorbid/Severe | to persons of normal weight. Therefore, these conditions are always clinically significant and reportable when documented by the provider ⁵ . |

For additional resources, contact our Provider Relations team at Providers@ARHealthWellness.com

Obesity & BMI Coding Tips



HEDIS*

Adult BMI Assessment

Measure Demonstrates percentage of members ages 18-74 who had an outpatient visit and whose body mass index (BMI) was documented during the measurement year or the year prior.

ICD-10 (age 20 or older):

Z68.1, Z68.20, Z68.21, Z68.22, Z68.23, Z68.24, Z68.25, Z68.26, Z68.27, Z68.28, Z68.29, Z68.30, Z68.31, Z68.32, Z68.33, Z68.34, Z68.35, Z68.36, Z68.37, Z68.38, Z68.39, Z68.41, Z68.42, Z68.43, Z68.44, Z68.45

Nutrition Counseling

ICD-10 Z71.3

CPT 97802, 97803, 97804

HCPCS G0270, G0271, G0447, S9449, S9452, S9470

Physical Activity Counseling

ICD-10 Z02.5, Z71.82

CPT 97802, 97803, 97804

HCPCS G0270, G0271, G0447, S9449, S9451, S9452, S9470

References and Notes

This educational material complies with accepted 2018 ICD-10-CM practices and guidelines and is for general supplemental purposes only. The information is not guaranteed to be complete, free of errors, or the most current revision. It is the responsibility of the provider to document accurate and complete codes, clinical rationale and medical services rendered to support appropriate ICD-10 code(s) according to official billing and coding guidelines, procedures, and regulations.

^{*}HEDIS measures can be found at www.ncga.com

¹ https://www.nhlbi.nih.gov/health-topics/overweight-and-obesity

²https://www.cms.gov/Medicare/Coding/ICD10/Downloads/2019-ICD10-Coding-Guidelines-.pdf

³Report on Medicare Compliance, Volume 26 Number 14 pg. 2. April 24, 2017

⁴https://www.nhlbi.nih.gov/health-topics/overweight-and-obesity

⁵AHA Coding Clinic, 3rd quarter 2011, Vol. 28, Num. 3, pages 4-5